

COMPARISION OF TWO EFFECTIVE ADVANCE PHYSIOTHERAPY INTERVENTION PROTOCOL IN UNILATERAL KNEE OSTEOARTHERITIS

AJEET YADAV¹, DR. NIDHI SHUKLA²

INTERNSHIP STUDENT¹, ASSISTANT PROFESSOR²

RAMA UNIVERSITY, KANPUR^{1,2}

ABSTRACT-

Osteoarthritis (OA), one of the top five most incapacitating conditions that affects more than one third of those over 65, is a degenerative joint disease characterised by joint inflammation and a reparative bone response. OA is a complete joint disease that affects joint ligaments, cartilage, menisci, and muscles that are connected to the joints due to various pathophysiological processes. It is essential to develop a treatment plan that take into account the physiology of degenerative joint disease, but it should also take into account the demands of the various age groups and populations.

The multi-structural physical, physiological, and functional deficits caused by the disease call for the development of a physiotherapy. The current study's goal is to assess the effectiveness of progressive resistance exercises, passive stretching exercises, soft tissue manipulation (MET), Maitland mobilisation, aerobic exercises, Tai chi, strength training, balneotherapy, and aquatic therapy that are planned, therapist-supervised, and patient-educated.

A convenience sample size of 100 was used for the initial investigation. The samples were divided into two study groups at random. BATS was the first interventional group, while PEPSMAN was the second. The interventional group's treatment plan included a physiotherapy routine that was monitored by a therapist.

The Visual Analogue scale, the modified WOMAC scale, the Time up and Go test, the functional reach test, the 40-meter fast-paced walk test, the stair climbing test, and the 30-second chair stand test were the outcome variables examined.

The majority of analysed end measures showed a significant improvement in the PEPSMAN interventional groups, and as a result, the planned supervised physiotherapy protocol was determined to be effective in treating the many physiological deficits linked to the whole joint illness.

KEY WORDS:

Maitland mobilisation, Progressive resistance exercise, Passive stretching exercise, Soft tissue manipulation, Balneotherapy, Muscle Energy Technique (MET), Knee OA (osteoarthritis), and PEPSMAN (Patient Education, Progressive Resistance Exercise, Muscle Energy Technique). BATS stands for balneotherapy, aqua exercise, tai chi, and strength training.

INTRODUCTION

The most prevalent articular illness, osteoarthritis, is projected to rank as the fourth most common cause of disability globally^[1]. However, of the entire disease burden, the knee joint accounts for 83% ^[2] of the disease burden in the majority of cases. Osteoarthritis of the knee joint has been shown to have considerable detrimental effects on quality of life, morbidity, and mortality, as well as the economic situation of the affected population ^[3]. A global prevalence of 16% was recorded in the population aged 15 and older in 2020, and

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG 22.9% of the population (654.1 million people) was 40 and older ^[4]. According to reports, 17–61% of people in India over the age of 50 have osteoarthritis of the knee joint ^[5]. The two most prominent issues that lead to mobility impairments, difficulties carrying out everyday tasks, and a poor quality of life in people with osteoarthritis of the knee joint are pain and reduced function ^[6]. The patients also had decreased proprioception, which can affect a person's postural stability and increase their risk of falling ^[7, 8, 9, 10].

Numerous research have offered numerous techniques, but none of them take a comprehensive approach. In one of the investigations, the population with knee osteoarthritis may have had symptom relief and an improvement in functional status with exercise therapy alone ^(13,14). Exercises for strengthening and cardio fitness were advised by the National Institute for Health and Care Excellence (NICE) ^(13,15). The osteoarthritis research society international (OARSI) recommended several different types of land-based structured exercise programmes, including balance, neuromuscular training, strengthening, cardio, and secondly, mind-body activities ^(13,16). Resistance workouts or aquatic aerobics were recommended by the American College of Rheumatology (ACR) ^(13,17). For the treatment of knee joints, the Ottawa panel recommended mind-body activities, aerobic exercises, and strengthening exercises with or without aerobic workouts ^(13,18,19,20). The EULAR advocated intensifying exercise and activity in general ^(13,21).

The goal of the current study is to offer patients with knee osteoarthritis an integrated physical therapy protocol that takes a comprehensive approach to managing physical and physiological deficits related to the various structures of the affected knee joint. The method we describe in this study is not only effective at preventing osteoarthritis, but it is also affordable and widely accessible to the general public.

MATERIALS AND METHODS

<u>STUDY DESIGN-</u>

The preliminary investigation, which included analysis according to protocol, included a convenience sample of 100 participants who were randomly assigned to the first interventional group (n = 50) and the second interventional group (n = 50) of two interventional groups. The names of all sixty participants were written on identical, well-shuffled chits, which were maintained in a jar to randomly assign participants to the first and second interventional groups. A person who was not involved in the study was requested to take one chit at a time out of the jar and place it into one or two bowls that were designated as the interventional groups, respectively. The Rama University in Kanpur, India approved the study and ensured that the two study groups were homogeneous. Before participating in the study, participants' patients provided written informed consent.

PARTICIPANTS-

Patients with knee osteoarthritis who were referred to Rama University's rehabilitation centre in Kanpur made up the patient sample pool. The 45–60 age range that was included, patients with a history of ligament and/or meniscal damage at the level of the knee joint, a recent fracture, cancer, another systemic disease, metabolic disease, infectious disease, or a history of neuromuscular disease, pregnant or nursing women, patients who had knee joint replacement surgery within the previous six months, or those who had received intra-articular injections of corticosteroids at the level of the knee joint, as well as those with known cases of ankylosing spondylitis, rheumatoid arthritis, osteomalacia, or Paget's diseases participants with a BMI of 35 or above and subjects with osteoarthritis worse than grade 2. Patients enrolled in or about to begin any other structured lower limb strengthening programme, including those without a formal structure, were all eliminated.

INTERVENTIONS-

The first interventional group's participants received the BATS physiotherapy programme, which includes balanotherapy, water treatment, Tai Chi, and strength training. The second interventional group underwent a therapist-supervised PEPSMAN physiotherapy protocol that included four sessions per week for six weeks of patient education, resistance training, passive stretching exercises, soft tissue manipulation, Maitland mobilisation, aerobic exercise, and neuromuscular training.

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG

Throughout the whole study period, the patients' education sessions were held once a week, for a total of four sessions. Each educational session was organised in such a way that the subsequent session was a progression of the previous one. Each one-on-one session lasted for an hour and a half. The meetings were held in a quiet, pleasant setting that was well-ventilated and free from outside distraction. The initial session's objectives were to inform the patient thoroughly about the sickness, comprehend his or her beliefs on the ailment, and assess the patient's health. In the first session, everything was explained in detail in terms that laypeople could understand, including what knee osteoarthritis is, its causes, related articular and periarticular changes, the need for treatment, the significance of precautions, the value of physiotherapy, and the function of our protocol. The second session's objectives were to encourage the patient and address any erroneous expectations, bad experiences, and misperceptions.

WEEKS	PEPSMAN GROUP	BATS GROUP
	Patient education,	Patient education,
FROM 1 WEEKS TO 2	progressive resistance	strength training.
WEEKS	e <mark>xer</mark> cises,	
	Passive stretching	Aquatic Therapy.
FROM 3 WEEKS TO 4	exercises, soft tissue	
WEEKS	manipulation.	
	Muscle energy	Tai chi Training.
FROM 4 WEEKS TO 5	technique, maitland	
WEEKS	mobilization.	
	Aerobic exercise,	Balneotherapy.
FROM 5 WEEK TO 6	neuromuscular training.	
WEEK		

International Research Journal Research Through Innovation

PROTOCOL



OUTCOMES

Pain intensity was one of the primary outcome measures evaluated. It was measured using the Visual Analogue Scale (VAS), which is a straight line that describes the Limits of pain, or the two extremes of pain—no pain at all and the worst suffering—are represented by the two ends of the line. The patient was asked to record the perceived pain level on the scale during the pre-test and post-test evaluation in order to account for the subjective nature of the pain. The difference between the two points on the scale that were reported by a person at two distinct times represented the actual variance in Pain intensity.

IJNRD2305705

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG In order to evaluate functional mobility, quality of life, and activities of daily living, the modified Western Ontario and McMaster University (WOMAC) Osteoarthritis Index was used.

During the pre-test and post-test evaluation phases, the participants received the index and were asked to report it as appropriate.

To evaluate the participants' dynamic steady state balance throughout the pre- and post-test periods,

The timed Up and Go test (TUG) was administered. To do this, it was timed how long it took the subject to stand up from a chair without armrests, walk five metres at a speed of his or her choosing, and then come back and sit down on the chair.

The objective of the functional reach test (FRT) was to evaluate dynamic steady state balance. The third metacarpal's initial position was marked on the participants after asking them to stand with their arms outstretched and fists made. The participants were then instructed to lean forward from the starting position without shifting their base of support until they reached the final position of maximum forward reach. For the aim of the assessment, the distance between the third metacarpal's knuckle's starting and final positions was measured.

The 40 metre fast paced walk test (40 m FPWT) was used to measure participants' ability to walk quickly over short distances, their ability to change directions while walking, and their speed while walking quickly. The participants were instructed to move swiftly and safely along a ten-meter pathway. At the end of the pathway, which was marked by a cone, participants were required to turn around and head back to the starting point, which was marked by another cone. This activity was repeated until the participants had successfully covered a forty-meter walkway without running. To gauge the pace of the walk, the amount of time from start to finish was recorded.

The stair climb test (SCT) evaluates a person's ability to climb or descend steps while simultaneously measuring the lower body's strength and balance. The participants were instructed to quickly and safely climb a flight of stairs with nine steps and an eight-inch step height at each, then turn and descend back to the ground, during the pre-test and post-test measurement periods [39]. Timing was taken from the starting position (both feet on the ground) to the ending position (both feet back on the ground).

The 30-second chair-stand test (30-s-CST) was used to evaluate the lower body's dynamic balance and strength while emphasising the subject's sit-to-stand action. The participants were instructed to cross their arms over their chest while sitting on a chair that was 17 inches high (seat height), with their knees bent at a 90-degree angle and their feet flat on the ground at shoulder width apart. The number of times the participants stood up and sat down during the thirty second exercise was recorded. The measurement was done both before and after the test.

Both clinically and statistically, PT has been proven to be significant. At 4 and 8 weeks, the treatment group demonstrated improvements in 6-min walk distance and WOMAC score, whereas the placebo group did not. Less discomfort, less joint stiffness, improved quality of life, and stronger hip muscles were the outcomes of these therapies. Approximately 72% and 75% of participants, vs only 17% (each) of control participants, reported improvements in pain and function, respectively.

RESULTS

Out of 80 samples recruited in the preliminary study, five patient from the 2nd interventional (BATS) group dropped out for a total knee replacement procedure during the study, the 1st interventional group experienced five dropouts due to unavoidable reasons, hence a total 35 samples in each 1st interventional group and 2nd interventional group completed the program. The characteristics of the samples are described in the exercises in 1st interventional group (PEPSMAN) physiotherapy protocol, consisting of patient education, progressive resistance exercise, passive stretching exercises and soft tissue manipulation, muscle energy technique, and maitland mobilization, aerobic exercises, and neuromuscular training. The second interventional (BATS) group are described the physiotherapy protocol –balneotherapy, aquatic, tai chi, and strength training exercises.

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG The overall significance for the chosen factor i.e., PEPSMAN and BATS, in the test of between sample effect and pairwise comparison was found to treatment. The administered interventional protocol consisting of a combination of exercise therapy and manual therapy is found significantly effective in the management of knee osteoarthritis patients, however in case of the stair climb and functional reach test,

The main data is adjusted for health, the 1st interventional group pre; and 1st interventional group post treatment. 2^{nd} interventional pre; and 2^{nd} interventional post treatment. The p < 0.001 and p < 0.01 is denoted. N= non significant.

DISCUSSION

A preliminary study is being conducted to assess the therapeutic potential of the therapist-supervised PEPSMAN physiotherapy regimen in a population of Indians with knee osteoarthritis to reduce pain and enhance static balance, dynamic balance, and physical function. Functional abilities, and hence, impairment. Evidence suggests that patient education is crucial in chronic illnesses like osteoarthritis, where patients first struggled to survive in a small space that discouraged them from engaging in healthy social contacts ^[13]. Disturbances in mental and physical health may be influenced by disturbed social wellbeing. Recent research has backed the use of appropriate patient education to aid in better illness management ^[13]. Exercise treatment is a well-researched and widely used therapeutic approach to treat knee osteoarthritis, but little research has been done to examine the therapeutic potential of manual therapy both alone and in combination with exercise therapy.

Patients with knee osteoarthritis were found to benefit significantly from an administered, designed, and therapist-supervised PEPSMAN physiotherapy protocol, which combines patient education, exercise therapy, and manual therapy. This protocol was found to have a significant therapeutic role in improving static balance, dynamic balance, pain, physical functional performance, and disability. The intervention data of the samples, which were acquired after four weeks, clearly show the outcome.

It is well known that ageing, which is a common cause of osteoarthritis, causes cell senescence, altered apoptosis, mitochondrial dysfunction, oxidative stress, and homeostatic dysfunction over time. As a result, the connective tissues that make up joints become stiff, fragile, and dehydrated. A delayed favorable response to the treatment in the older age group, compared to the younger ones, may be due to the continuum of harmful alterations brought on by ageing.

An evaluation of balance is the functional reach test. Although the outcome measure of balance is frequently used in situations with lower limb deficits, the ability to maintain balance is a combined function of the lower limb and trunk stabilisers. Despite the fact that both groups experienced pain alleviation, the difference in pain between the two groups was not very noticeable. Separately, manual treatment and exercise therapy both showed promise in assisting with the disease's symptom reduction.

Being a chronic ailment, knee osteoarthritis incorporates a psychological aspect of pain. The presented non significant difference between the two groups, examined at the end of the trial, may be due to subjective belief and satisfaction.

In comparison to the BATS group, the intervention group's static and dynamic balance dramatically improved. Muscle strength, flexibility, conditioning, and joint proprioception were among the parameters that were to be improved through the physiotherapeutic exercises and therapy given to the intervention group; as a result, it is possible that the improvement in these parameters led to an improvement in the samples' relative balance.

As described in the research, individuals with knee OA frequently combine a non-pharmacological exercise and therapy routine with pharmacological treatment options to achieve better results.^[9,10,12] Despite the therapeutic options available, a number of deficiencies continue to exist, making cost containment a serious concern.^[15] Approximately half of all OA patients undergoing physiotherapy are still using subpar diagnostic techniques, recommendations, and treatment modalities that lack proof of benefit (including massage, traction, and stretching).^[15]

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG Given that there are now no adequate and effective treatment regimens, this presents a serious healthcare challenge for persons who have OA-related issues. As can be shown above with only one ailment, the future demands for healthcare vastly exceed the current ways of approach. Therefore, in order to define the diagnostic processes, assessments, treatment modalities, and evaluations required for a future programe that is more cost- and treatment-effective, higher standards of quality research will be essential.

LIMITATIONS OF THE STUDY

In this study paced dropouts as was unable to follow up. The sample size and duration of the study is also reduce.

RECOMMENDATIONS FOR FUTURE STUDIES

For Further study with a similar approach will be conducted with more advance techniques on a large sample size with increase study duration.

REFRENCE

- Murray C.J., Vos T., Lozano R., Naghavi M., Flaxman A.D., Michaud C., Ezzati M., Shibuya K., Salomon J.A., Abdalla S., et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. Lancet. 2012;380:2197–2223. doi: 10.1016/S0140-6736(12)61689-4. - DOI - PubMed
- Vos T., Flaxman A.D., Naghavi M., Lozano R., Michaud C., Ezzati M., Shibuya K., Salomon J.A., Abdalla S., Aboyans V., et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. Lancet. 2012;380:2163–2196. doi: 10.1016/S0140-6736(12)61729-2. - <u>DOI</u> - <u>PMC</u> - <u>PubMed</u>
- Briggs A.M., Cross M.J., Hoy D.G., Sanchez-Riera L., Blyth F.M., Woolf A.D., March L. Musculoskeletal health conditions represent a global threat to healthy aging: A report for the 2015 World Health Organization world report on ageing and health. Gerontologist. 2016;56((Suppl. 2)):S243–S255. doi: 10.1093/geront/gnw002. - <u>DOI</u> - <u>PubMed</u>
- Cui A., Li H., Wang D., Zhong J., Chen Y., Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. EClinicalMedicine. 2020;29:100587. doi: 10.1016/j.eclinm.2020.100587. - <u>DOI</u> - <u>PMC</u> - <u>PubMed</u>
- Kaur R., Sharma V.L., Singh A. Prevalence of knee osteoarthritis and its correlation in women of rural and urban parts of Hoshiarpur (Punjab) J. Postgrad. Med. Educ. Res. 2015;49:32. doi: 10.5005/jpjournals-10028-1139. - DOI
- Odole A.C., Ogunlana M.O., Adegoke B.O., Ojonima F., Useh U. Depression, pain and physical function in patients with osteoarthritis of the knee: Implications for interprofessional care. Niger. J. Med. Rehabil. 2015;18 doi: 10.34058/njmr.v18i1.120. - DOI
- Knoop J., Steultjens M.P., Van der Leeden M., Van der Esch M., Thorstensson C.A., Roorda L.D., Lems W.F., Dekker J. Proprioception in knee osteoarthritis: A narrative review. Osteoarthr. Cartil. 2011;19:381–388. doi: 10.1016/j.joca.2011.01.003. - <u>DOI</u> - <u>PubMed</u>
- Bascuas I., Tejero M., Monleón S., Boza R., Muniesa J.M., Belmonte R. Balance 1 year after TKA: Correlation with clinical variables. Orthopedics. 2013;36:e6–e12. doi: 10.3928/01477447-20121217-11. - DOI - PubMed
 - a. Ringdahl E, Pandit S. Treatment of knee osteoarthritis. *Am Fam Physician*. 2011;83:1287–92. [PubMed] [Google Scholar]

- © 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG 9. Sharma L., Pai Y.C., Holtkamp K., Rymer W.Z. Is knee joint proprioception worse in the arthritic knee versus the unaffected knee in unilateral knee osteoarthritis? Arthritis Rheum. Off. J. Am. Coll. Rheumatol. 1997;40:1518–1525. doi: 10.1002/art.1780400821. - <u>DOI</u> - <u>PubMed</u> Magrans-Courtney T, Wilborn C, Rasmussen C, Ferreira M, Greenwood L, Campbell B, et al. Effects of diet type and supplementation of glucosamine, chondroitin, and MSM on body composition, functional status, and markers of health in women with knee osteoarthritis initiating a resistance-based exercise and weight loss program. *J Int Soc Sports Nutr.* 2011;8:8. [PMC free article] [PubMed] [Google Scholar]
- Khalaj N., Abu Osman N.A., Mokhtar A.H., Mehdikhani M., Wan Abas W.A. Balance and risk of fall in individuals with bilateral mild and moderate knee osteoarthritis. PLoS ONE. 2014;9:e92270. doi: 10.1371/journal.pone.0092270. - <u>DOI</u> - <u>PMC</u> - <u>PubMed</u>

a. Esser S, Bailey A. Effects of exercise and physical activity on knee OA. *Curr Pain Headache Rep.* 2011;15:423–30. [PubMed] [Google Scholar

- Dantas L.O., de Fátima Salvini T., McAlindon T.E. Knee osteoarthritis: Key treatments and implications for physical therapy. Braz. J. Phys. Ther. 2021;25:135–146. doi: 10.1016/j.bjpt.2020.08.004. - DOI - PMC - PubMed
- Fransen M., McConnell S., Harmer A.R., Van der Esch M., Simic M., Bennell K.L. Exercise for osteoarthritis of the knee. Cochrane Database Syst. Rev. 2015;1:CD004376. doi: 10.1002/14651858.CD004376.pub3. - <u>DOI</u> - <u>PubMed</u>
 - a. Kon E, Filardo G, Drobnic M, Madry H, Jelic M, Dijk N, et al. Non-surgical management of early knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc.* 2012;20:436–9. [PubMed] [Google Scholar]
- National Clinical Guideline Centre (UK) Osteoarthritis: Care and Management in Adults. National Institute for Health and Care Excellence; London, UK: 2014. - <u>PubMed</u>
- Bannuru R.R., Osani M.C., Vaysbrot E.E., Arden N.K., Bennell K., Bierma-Zeinstra S.M., Kraus V.B., Lohmander L.S., Abbott J.H., Bhandari M., et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. Osteoarthr. Cartil. 2019;27:1578–1589. doi: 10.1016/j.joca.2019.06.011. - <u>DOI</u> - <u>PubMed</u>
- 15. Kolasinski S.L., Neogi T., Hochberg M.C., Oatis C., Guyatt G., Block J., Callahan L., Copenhaver C., Dodge C., Felson D., et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. Arthritis Rheumatol. 2020;72:220-233. doi: 10.1002/art.41142. - DOI - PubMed Smink AJ, van den Ende CH, Vliet Vlieland TP. Beating osteoarthritis: Development of a stepped care strategy to optimize utilization and timing of nontreatment modalities for patients with hip or knee osteoarthritis. Clin surgical Rheumatol. 2011;30:1623–9. [PubMed] [Google Scholar]
- 16. Brosseau L., Taki J., Desjardins B., Thevenot O., Fransen M., Wells G.A., Imoto A.M., Toupin-April K., Westby M., Gallardo I.C., et al. The Ottawa panel clinical practice guidelines for the management of knee osteoarthritis. Part one: Introduction, and mind-body exercise programs. Clin. Rehabil. 2017;31:582–595. doi: 10.1177/0269215517691083. DOI PubMed
- Brosseau L., Taki J., Desjardins B., Thevenot O., Fransen M., Wells G.A., Mizusaki Imoto A., Toupin-April K., Westby M., Alvarez Gallardo I.C., et al. The Ottawa panel clinical practice guidelines for the management of knee osteoarthritis. Part two: Strengthening exercise programs. Clin. Rehabil. 2017;31:596–611. doi: 10.1177/0269215517691084. - <u>DOI</u> - <u>PubMed</u>
- 18. Brosseau L., Taki J., Desjardins B., Thevenot O., Fransen M., Wells G.A., Mizusaki Imoto A., Toupin-April K., Westby M., Alvarez Gallardo I.C., et al. The Ottawa panel clinical practice guidelines for the

© 2023 IJNRD | Volume 8, Issue 5 May 2023 | ISSN: 2456-4184 | IJNRD.ORG management of knee osteoarthritis. Part three: Aerobic exercise programs. Clin. Rehabil. 2017;31:612–624. doi: 10.1177/0269215517691085. - <u>DOI</u> - <u>PubMed</u>

- Fernandes L., Hagen K.B., Bijlsma J.W., Andreassen O., Christensen P., Conaghan P.G., Doherty M., Geenen R., Hammond A., Kjeken I., et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. Ann. Rheum. Dis. 2013;72:1125–1135. doi: 10.1136/annrheumdis-2012-202745. - DOI - PubMed
- 20. Glossary. [(accessed on 27 November 2022)];Spine. 2000 25:3200–3202. doi: 10.1097/00007632-200012150-00016. Available online: <u>https://journals.lww.com/spinejournal/fulltext/2000/12150/glossary.16.aspx</u>. - <u>DOI</u>
- 21. Haefeli M., Elfering A. Pain assessment. Eur. Spine J. 2006;15:S17–S24. doi: 10.1007/s00586-005-1044-x. - DOI - PMC - PubMed
- 22. Price D.D., Harkins S.W., Baker C. Sensory-affective relationships among different types of clinical and experimental pain. Pain. 1987;28:297–307. doi: 10.1016/0304-3959(87)90065-0.
 <u>DOI</u> <u>PubMed</u>
- Price D.D., McGrath P.A., Rafii A., Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. Pain. 1983;17:45–56. doi: 10.1016/0304-3959(83)90126-4. - <u>DOI</u> - <u>PubMed</u>
- 24. Samuel A.J., Kanimozhi D. Outcome measures used in patient with knee osteoarthritis: With special importance on functional outcome measures. Int. J. Health Sci. 2019;13:52. <u>PMC</u> <u>PubMed</u>
- 25. Bergquist R., Weber M., Schwenk M., Ulseth S., Helbostad J.L., Vereijken B., Taraldsen K. Performance-based clinical tests of balance and muscle strength used in young seniors: A systematic literature review. BMC Geriatr. 2019;19:9. doi: 10.1186/s12877-018-1011-0. DOI PMC PubMed
- 26. Dobson F., Hinman R.S., Roos E.M., Abbott J.H., Stratford P., Davis A.M., Buchbinder R., Snyder-Mackler L., Henrotin Y., Thumboo J., et al. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. Osteoarthr. Cartil. 2013;21:1042–1052. doi: 10.1016/j.joca.2013.05.002. DOI PubMed
- 27. Maly M.R., Krupa T. Personal experience of living with knee osteoarthritis among older adults. Disabil. Rehabil. 2007;29:1423–1433. doi: 10.1080/09638280601029985. <u>DOI</u> <u>PubMed</u>
- 28. Nyvang J., Hedström M., Gleissman S.A. It's not just a knee, but a whole life: A qualitative descriptive study on patients' experiences of living with knee osteoarthritis and their expectations for knee arthroplasty. Int. J. Qual. Stud. Health Well-Being. 2016;11:30193. doi: 10.3402/qhw.v11.30193.
 <u>DOI</u> <u>PubMed</u>
- 29. Pouli N., Das Nair R., Lincoln N.B., Walsh D. The experience of living with knee osteoarthritis: Exploring illness and treatment beliefs through thematic analysis. Disabil. Rehabil. 2014;36:600–607. doi: 10.3109/09638288.2013.805257. - DOI - PMC - PubMed Deyle GD, Allison SC, Matekel RL, Ryder MG, Stang JM, Gohdes DD, et al. Physical therapy treatment effectiveness for osteoarthritis of the knee: A randomized comparison of supervised clinical exercise and manual therapy procedures versus a home exercise program. *Phys Ther*. 2005;85:1301–17. [PubMed] [Google Scholar]
- 30. Lo GH, LaValley M, McAlindon T, Felson D. Intra-articular hyaluronic acid in treatment of knee osteoarthritis. *JAMA*. 2003;290:3115–21. [PubMed] [Google Scholar]