

# Anti-Inflammatory Potential of *Bryophyllum pinnatum*: A Review of Phytochemistry, Pharmacology, and Therapeutic Applications

Ms Pramila Suresh Chaugule<sup>1</sup>, Dr. A.R. Aswar<sup>2</sup>, Dr. Shrikant M Darekar<sup>3</sup>,  
Dr. Manisha Zaware<sup>4</sup>

Research scholar<sup>1</sup>, Research Guide<sup>2,3,4</sup>

H.S.B.P.V.T'S GOI Faculty of pharmacy, Kashti<sup>1,2,3,4</sup>

## Abstract

Inflammation is a complex biological response of the body to harmful stimuli such as pathogens, tissue injury, toxins, and autoimmune reactions. Although inflammation serves as a protective mechanism, prolonged or excessive inflammatory responses contribute to the development of various chronic diseases, including arthritis, cardiovascular disorders, asthma, and inflammatory bowel disease. Conventional anti-inflammatory drugs such as non-steroidal anti-inflammatory drugs (NSAIDs) and corticosteroids are widely used for the management of inflammatory conditions; however, their long-term use is often associated with adverse effects including gastrointestinal irritation, renal toxicity, cardiovascular complications, and immunosuppression. These limitations have stimulated interest in the exploration of safer and more effective plant-based therapeutic agents.

*Bryophyllum pinnatum* (Lam.) Oken, a medicinal plant belonging to the family Crassulaceae, has been extensively utilized in traditional medicine for the treatment of inflammation, pain, wounds, infections, and various other ailments. The plant contains numerous bioactive phytoconstituents, including flavonoids, phenolic compounds, tannins, alkaloids, glycosides, saponins, steroids, and triterpenoids, which are reported to possess significant pharmacological activities. Several experimental studies have demonstrated the anti-inflammatory potential of *Bryophyllum pinnatum* using models such as carrageenan-induced paw edema, formaldehyde-induced inflammation, and cotton pellet granuloma. The observed effects are attributed to the inhibition of inflammatory mediators, suppression of pro-inflammatory cytokines, antioxidant activity, and modulation of inflammatory signaling pathways.

The available scientific evidence supports the traditional use of *Bryophyllum pinnatum* as a natural anti-inflammatory agent. Further studies focusing on isolation of active constituents, standardization of extracts, elucidation of molecular mechanisms, and clinical evaluation are necessary to establish its therapeutic efficacy and promote its development as a potential phytopharmaceutical for inflammatory disorders.

**Keywords:** *Bryophyllum pinnatum*, Inflammation, Anti-inflammatory activity, Medicinal plants, Phytochemicals, Flavonoids, Cytokines, Herbal medicine, Carrageenan-induced paw edema, Phytopharmaceuticals.

## Introduction

### Inflammation

Inflammation is a protective biological response of the body against harmful stimuli such as infections, injuries, toxins, and tissue damage. It is characterized by redness, swelling, heat, pain, and loss of function at the affected site. Inflammation helps eliminate harmful agents and initiates tissue repair. Acute inflammation is a short-term response that develops rapidly and resolves within a few days, whereas chronic inflammation persists for a prolonged period and may contribute to diseases such as arthritis, cardiovascular disorders, diabetes, and inflammatory bowel disease. Various inflammatory mediators, including histamine, prostaglandins, and cytokines (TNF- $\alpha$ , IL-1 $\beta$ , and IL-6), play crucial roles in regulating and amplifying the inflammatory response.

### Need for Herbal Anti-inflammatory Agents

Conventional anti-inflammatory drugs such as non-steroidal anti-inflammatory drugs (NSAIDs) and corticosteroids are commonly used to manage inflammatory conditions. However, prolonged use of these medications may lead to adverse effects including gastric irritation, peptic ulcers, renal impairment, cardiovascular complications, osteoporosis, and immunosuppression. These limitations have increased the demand for safer and more effective alternatives. Medicinal plants have gained considerable attention due to their natural origin, therapeutic potential, lower incidence of side effects, and long history of traditional use.

### Importance of *Bryophyllum pinnatum*

*Bryophyllum pinnatum* (Lam.) Oken is a medicinal plant belonging to the family Crassulaceae and is widely used in traditional medicine for the treatment of inflammation, pain, wounds, ulcers, infections, and various other ailments. The plant is rich in bioactive compounds such as flavonoids, phenolics, tannins, glycosides, and triterpenoids, which possess anti-inflammatory and antioxidant properties. Its extensive ethnomedicinal use and promising pharmacological activities make it an important candidate for scientific investigation. Therefore, *Bryophyllum pinnatum* has been selected for anti-inflammatory studies to validate its traditional claims and explore its potential as a natural therapeutic agent.

### Inflammation and Anti-inflammatory Mechanisms

#### Inflammation

Inflammation is a complex protective response of the body's immune system to harmful stimuli such as infections, tissue injury, toxins, or irritants. Its primary function is to eliminate the causative agent, remove damaged tissues, and initiate the healing process. Although inflammation is essential for host defense, excessive or prolonged inflammation can lead to tissue damage and various chronic diseases.

The classical signs and symptoms of inflammation include redness (rubor), heat (calor), swelling (tumor), pain (dolor), and loss of function (functio laesa). These manifestations result from increased blood flow, enhanced vascular permeability, and migration of immune cells to the site of injury.

The inflammatory response begins when tissue injury or infection activates immune cells such as macrophages and mast cells. These cells release various inflammatory mediators, including histamine, prostaglandins, leukotrienes, and cytokines, which promote vasodilation, increase vascular permeability, and attract leukocytes to the affected area. The recruited inflammatory cells remove pathogens and damaged tissue

through phagocytosis and release additional mediators that amplify the response. Once the harmful stimulus is eliminated, anti-inflammatory mechanisms facilitate tissue repair and restore normal physiological function. However, failure to resolve inflammation may result in chronic inflammatory conditions such as arthritis, asthma, cardiovascular diseases, and inflammatory bowel disorders.

### Mechanism of Anti-inflammatory Action

Anti-inflammatory agents exert their effects through multiple mechanisms that interfere with the production and action of inflammatory mediators. One of the major mechanisms involves inhibition of the cyclooxygenase (COX) pathway, which reduces the synthesis of prostaglandins responsible for pain, swelling, and inflammation. Suppression of inflammatory cytokines such as tumor necrosis factor-alpha (TNF- $\alpha$ ), interleukin-1 beta (IL-1 $\beta$ ), and interleukin-6 (IL-6) also plays an important role in controlling inflammatory responses and preventing tissue damage.

Many medicinal plants possess antioxidant properties that help reduce oxidative stress by scavenging reactive oxygen species (ROS) and free radicals generated during inflammation. Reduction of oxidative stress prevents cellular injury and limits the progression of inflammatory disorders. Another important mechanism involves modulation of Nuclear Factor-kappa B (NF- $\kappa$ B), a key transcription factor that regulates the expression of numerous inflammatory genes. Inhibition of NF- $\kappa$ B activation decreases the production of cytokines, chemokines, and other inflammatory mediators, thereby attenuating the inflammatory response.

The anti-inflammatory activity of *Bryophyllum pinnatum* is believed to be associated with these mechanisms due to the presence of bioactive phytoconstituents such as flavonoids, phenolic compounds, tannins, and triterpenoids, which collectively contribute to its therapeutic potential.

### Plant Profile of *Bryophyllum pinnatum*

#### Taxonomical Classification

Category	Classification
Kingdom	Plantae
Division	Magnoliophyta
Class	Magnoliopsida
Order	Saxifragales
Family	Crassulaceae
Genus	Bryophyllum
Species	<i>Bryophyllum pinnatum</i>

#### Common Names

- Life Plant
- Miracle Leaf
- Air Plant
- Cathedral Bells
- Patharchatta (India)

## Botanical Description

- Succulent perennial herb.
- Thick fleshy leaves.
- Opposite leaves with serrated margins.
- Bell-shaped flowers.

## Geographical Distribution

- India
- Africa
- South America
- Southeast Asia

## Phytochemical Constituents

- Flavonoids
- Phenolic compounds
- Tannins
- Alkaloids
- Glycosides
- Triterpenoids
- Steroids
- Saponins

## Traditional Uses

- Anti-inflammatory
- Wound healing
- Antimicrobial
- Antidiabetic
- Analgesic
- Antiulcer

## Experimental Studies

### Carrageenan-Induced Paw Edema

The carrageenan-induced paw edema model is one of the most widely used experimental methods for evaluating acute anti-inflammatory activity. In this model, carrageenan is injected into the hind paw of experimental animals, producing localized inflammation characterized by edema formation. Several studies have reported that *Bryophyllum pinnatum* extract significantly reduces paw swelling, indicating its ability to inhibit inflammatory mediators involved in the acute phase of inflammation.

### Formaldehyde-Induced Inflammation

The formaldehyde-induced paw edema model is used to assess sub-acute and chronic inflammatory responses. Injection of formaldehyde into the paw produces persistent inflammation resembling arthritic conditions.

Treatment with *Bryophyllum pinnatum* extract has been shown to reduce paw edema and inflammatory symptoms, suggesting its effectiveness in controlling prolonged inflammatory reactions and tissue damage.

### Cotton Pellet Granuloma Model

The cotton pellet granuloma model is commonly employed to evaluate the proliferative phase of chronic inflammation. In this method, sterile cotton pellets are implanted subcutaneously, leading to granuloma formation due to fibroblast proliferation and collagen deposition. Studies have demonstrated that *Bryophyllum pinnatum* extract significantly decreases granuloma weight, indicating its ability to suppress chronic inflammatory processes and connective tissue proliferation.

### Reported Findings

Various pharmacological studies have demonstrated the significant anti-inflammatory potential of *Bryophyllum pinnatum*. Administration of the plant extract has been reported to produce a marked reduction in edema formation in carrageenan-induced and formaldehyde-induced inflammation models, indicating its effectiveness against both acute and chronic inflammatory conditions.

The anti-inflammatory activity of *Bryophyllum pinnatum* is attributed to its ability to inhibit the production and release of inflammatory mediators such as prostaglandins, histamine, leukotrienes, and pro-inflammatory cytokines including TNF- $\alpha$ , IL-1 $\beta$ , and IL-6. This inhibition helps reduce swelling, pain, and tissue damage associated with inflammation.

In addition, the plant exhibits considerable antioxidant activity due to the presence of flavonoids, phenolic compounds, and other bioactive constituents. These compounds scavenge reactive oxygen species and free radicals, thereby reducing oxidative stress and preventing cellular injury during inflammatory processes.

Histopathological studies further support the anti-inflammatory effects of *Bryophyllum pinnatum*. Microscopic examination of inflamed tissues has shown reduced edema, decreased inflammatory cell infiltration, lower vascular congestion, and improved tissue architecture in extract-treated groups compared to untreated controls. These findings collectively validate the traditional use of *Bryophyllum pinnatum* as a potential natural anti-inflammatory agent.

### Mechanisms Proposed

The anti-inflammatory activity of *Bryophyllum pinnatum* is believed to result from multiple pharmacological mechanisms. One of the primary mechanisms involves inhibition of the cyclooxygenase (COX) pathway, which reduces the synthesis of prostaglandins responsible for inflammation, pain, and edema. By decreasing prostaglandin production, the plant extract helps alleviate inflammatory symptoms.

Another important mechanism is the suppression of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF- $\alpha$ ), interleukin-1 beta (IL-1 $\beta$ ), and interleukin-6 (IL-6). Reduction in the levels of these cytokines limits the recruitment and activation of inflammatory cells, thereby attenuating the inflammatory response.

*Bryophyllum pinnatum* also exhibits significant antioxidant activity due to the presence of flavonoids, phenolic compounds, tannins, and triterpenoids. These phytoconstituents neutralize reactive oxygen species (ROS) and free radicals generated during inflammation, reducing oxidative stress and protecting tissues from damage.

In addition, the plant extract may stabilize cellular and lysosomal membranes, preventing the release of inflammatory enzymes and mediators into surrounding tissues. This membrane-stabilizing effect contributes to the reduction of tissue injury and further supports its anti-inflammatory action. The combined effects of COX inhibition, cytokine suppression, antioxidant activity, and membrane stabilization account for the therapeutic potential of *Bryophyllum pinnatum* in inflammatory disorders.

## Future Perspectives

Despite the promising anti-inflammatory potential of *Bryophyllum pinnatum*, further research is required to fully establish its therapeutic value. Isolation and characterization of the specific bioactive phytoconstituents responsible for its anti-inflammatory activity will help in understanding its mechanism of action and facilitate the development of novel plant-based drugs.

Standardization of plant extracts is essential to ensure consistent quality, safety, efficacy, and reproducibility of therapeutic effects. Future studies should focus on developing standardized extraction procedures and quality control parameters for herbal preparations of *Bryophyllum pinnatum*.

The formulation of herbal products such as tablets, capsules, gels, ointments, and nanoformulations may improve the stability, bioavailability, and patient acceptability of the plant extract. Such formulations could enhance its therapeutic applications in the management of inflammatory conditions.

Comprehensive clinical trials and long-term safety evaluations are necessary to validate the efficacy and safety of *Bryophyllum pinnatum* in humans. These studies will provide scientific evidence to support its use in modern healthcare systems.

Furthermore, *Bryophyllum pinnatum* has the potential to be used as an adjunct therapy alongside conventional anti-inflammatory medications. Its natural origin, antioxidant properties, and anti-inflammatory effects may help improve treatment outcomes while reducing the dosage and adverse effects associated with synthetic drugs.

## Conclusion

*Bryophyllum pinnatum* is an important medicinal plant that has gained considerable attention due to its significant anti-inflammatory potential. The plant has been widely used in traditional medicine for the treatment of inflammation, pain, wounds, and various other ailments. Scientific investigations have demonstrated that its pharmacological activity is largely attributed to the presence of bioactive phytoconstituents such as flavonoids, phenolic compounds, tannins, and triterpenoids, which possess anti-inflammatory and antioxidant properties.

Various experimental studies, including carrageenan-induced paw edema, formaldehyde-induced inflammation, and cotton pellet granuloma models, have confirmed the ability of *Bryophyllum pinnatum* extracts to reduce inflammation and associated tissue damage. These findings provide scientific support for its traditional use in the management of inflammatory disorders.

The plant represents a promising natural alternative to conventional anti-inflammatory drugs, which are often associated with adverse effects during long-term use. However, further research is necessary to isolate and characterize its active constituents, standardize herbal preparations, and clarify the underlying molecular mechanisms responsible for its therapeutic effects. Well-designed clinical studies are also required to establish its efficacy, safety, dosage, and potential therapeutic applications in human inflammatory diseases. Overall,

*Bryophyllum pinnatum* holds considerable promise as a future phytopharmaceutical agent for the treatment and management of inflammatory conditions.

## References

1. Pal SK, Sen T, Chaudhuri AK. Neuropsychopharmacological profile of the methanolic fraction of *Bryophyllum pinnatum* leaf extract. *J Pharm Pharmacol*. 1999;51(3):313-318.
2. Pal S, Sen T, Chaudhuri AK. Analgesic and anti-inflammatory activities of the methanolic extract of *Bryophyllum pinnatum* leaves. *J Ethnopharmacol*. 1999;66(2):195-201.
3. Tanko Y, Mohammed A, Okasha MA, Umar AH, Magaji RA. Anti-nociceptive and anti-inflammatory activities of ethanol extract of *Bryophyllum pinnatum* leaves in experimental animals. *Afr J Biotechnol*. 2008;7(18):3363-3367.
4. Onwubuya EI, Chikezie PC, Uwakwe AA. Anti-inflammatory activity of hydroethanolic leaf extract of *Bryophyllum pinnatum* in carrageenan-induced paw edema and cotton pellet granuloma models. *Int J Pharm Sci Res*. 2021;12(4):2100-2108.
5. Oladejo AA, Chikezie PC, Uwakwe AA. Bioactive compounds and anti-inflammatory effects of hydroethanolic leaf extract of *Bryophyllum pinnatum* in experimental rats. *J Complement Integr Med*. 2022;19(3):557-566.
6. Ogunmoyole T, Olalekan OO, Ojo OA. Antioxidant and antidiabetic properties of *Bryophyllum pinnatum* leaf extract and identification of bioactive compounds using GC-MS. *Heliyon*. 2023;9(4):e15032.
7. Handique C, Dutta A, Saikia D. Evaluation of antinociceptive activity of ethanolic extract of *Bryophyllum pinnatum* leaves in mice. *Int J Basic Clin Pharmacol*. 2015;4(2):268-272.
8. Pandurangan A, Khosa RL, Hemalatha S. Evaluation of anti-inflammatory activity of *Bryophyllum calycinum* using carrageenan-induced paw edema and cotton pellet granuloma models. *Int J Pharm Pharm Sci*. 2011;3(3):125-128.
9. Chaithra SR, Nagesh KS. A review on phytochemistry and pharmacological activities of *Bryophyllum pinnatum*. *Int J Res Pharm Sci*. 2020;11(4):6840-6848.
10. Ukamaka E, Nwafor SV, Okeke CU. Phytochemistry, pharmacological activities and therapeutic potential of *Bryophyllum pinnatum*: A review. *Int J Pharm Sci Rev Res*. 2024;84(1):45-58.
11. Kamboj A, Saluja AK. *Bryophyllum pinnatum* (Lam.) Kurz.: Phytochemical and pharmacological profile. *Pharmacogn Rev*. 2009;3(6):364-374.
12. Yadav NP, Dixit VK. Recent approaches in herbal drug standardization and quality control. *Int J Integr Biol*. 2008;2(3):195-203.
13. Gupta VK, Sharma SK. Plants as natural anti-inflammatory agents: A review. *Int J Green Pharm*. 2017;11(2):S89-S98.
14. Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008;454(7203):428-435.
15. Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*. 2018;9(6):7204-7218.
16. Libby P. Inflammatory mechanisms: The molecular basis of inflammation and disease. *Nutr Rev*. 2007;65(12 Pt 2):S140-S146.
17. Ricciotti E, FitzGerald GA. Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol*. 2011;31(5):986-1000.
18. Lawrence T. The nuclear factor NF- $\kappa$ B pathway in inflammation. *Cold Spring Harb Perspect Biol*. 2009;1(6):a001651.
19. Harborne JB. *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. 3rd ed. London: Chapman and Hall; 1998.
20. Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 56th ed. Pune: Nirali Prakashan; 2021.