


MANAGEMENT OF 1ST STAGE CANCER WITHOUT MEDICATION

Malleswari.K, Dr.Rama Brahma Reddy.D, Bhavitha.M

Pharmaceutics, Phytochemistry, Student of pharmacy
Nalanda institute of pharmaceutical sciences



ABSTRACT

Cancer remains a major global health challenge, accounting for one in six deaths worldwide. Conventional treatments such as surgery, chemotherapy, and radiation are effective but often associated with significant side effects. In contrast, nonpharmacological approaches—including meditation, yoga, hypnosis, acupuncture, aromatherapy, music therapy, massage, and tai chi—offer supportive benefits in the management of Stage 1 cancer. These integrative therapies, while not replacements for medical treatment, can reduce stress, anxiety, and fatigue, improve sleep and mood, and enhance overall quality of life. Early detection of Stage 1 cancer provides a critical opportunity for intervention, and combining lifestyle modifications with supportive therapies can help patients maintain resilience and wellbeing during their cancer journey.

KEYWORD

Cancer, stage 1 cancer Early detection Nonpharmacological treatment integrative medicine alternative therapy macological treatment

INTRODUCTION

Cancer is one of the leading global health challenges, responsible for nearly one in six deaths worldwide. In 2020 alone, an estimated 19.3 million new cases and 10 million deaths were reported, highlighting its immense burden on public health. Cancer is not a single disease, but a complex group of conditions characterized by uncontrolled cell growth and gradual progression from localized tumors to widespread metastasis.

For decades, conventional treatments such as surgery, radiation therapy, and chemotherapy have been the primary options for patients. While these approaches remain central to cancer management, they often involve significant side effects and complications. In recent years, advances in targeted therapies, biological molecules, and immune-mediated treatments have expanded the possibilities for care.

At the same time, integrative and non-pharmacological methods—such as meditation, yoga, hypnosis, acupuncture, aromatherapy, and music therapy—have gained attention for their supportive role in managing early-stage cancers. These approaches, though not replacements for medical treatment, can reduce stress, improve quality of life, and help patients cope with the challenges of cancer therapy.

Thus, the management of Stage 1 cancer emphasizes the importance of early detection, lifestyle modification, and complementary therapies that enhance resilience and recovery alongside medical guidance




FIG:1Cancer cell




FIG 2: Stages of cancer

EARLY DETECTION

The greatest advantage when it comes to cancer treatment is the timely detection. In Stage 1 cancers, the disease is usually localised and has not spread to lymph nodes or distant parts of the body. The five-year survival rates for Stage 1 cancers, like breast, prostate, skin (melanoma), and thyroid, are significantly higher compared to later stages.

TYPES OF CANCER

There are over 100 types of cancer. Healthcare providers categorize them according to where they start in your body and the type of tissue they affect. There are three broad cancer classifications:

Solid cancers:

This is the most common type of cancer, making up about 80% to 90% of all cases. This includes carcinoma that forms in epithelial tissue (like your skin, breast, colon and lungs) and sarcoma that forms in bone and connective tissues.

Blood cancers:

These are cancers that start in your blood cells or lymphatic system. Examples include leukemia, lymphoma and multiple myeloma. Mixed: Cancers that involve two classifications or subtypes. Examples include carcinosarcoma and adenosquamous carcinoma.




FIG:3Types of cancer

Carcinomas cancer

Sarcomas cancer

Melanoma cancer

Leukemias cancer

Lymphomas cancer

ETIOLOGY OF CANCER

Modeling human situations such as lack of motivation to perform, comorbid conditions, and concomitant medications cannot yet be recapitulated. Valid preclinical models can the causes and contributing factors of fatigue are undoubtedly multifactorial, and thus its etiology is not well delineated. Understanding the causative factors is important for finding effective clinical therapies or preventative strategies for CRF. Patients with cancer experience multiple comorbid conditions, in addition to the malignancy itself, that can cause or exacerbate fatigue. This combination of factors may prevent or delay detection of the ultimate contributing disorders. Patients may acquire these medical conditions after receiving therapy or they may have these conditions at the time of presentation with cancer; the term “cancer-related” can encompass all these factors, because the exact time of onset in the cancer trajectory may be unknown. Defining CRF is quite challenging because many patients with cancer present with fatigue before primary adjuvant therapy is begun. Regardless of the source, clinically recognizing separate causes of fatigue at presentation will help uncover treatable situations, such as anemia, hypothyroidism, depression, concomitant medications, and infection. Remediating potentially reversible conditions is the initial approach to CRF according to NCCN guidelines.

The literature regarding possible CRF mechanisms contains a few small clinical trials and little preclinical work. The published data suggest multiple nonspecific interactions between cancer, the effects from cancer therapy, and the patients' responses to both. Unfortunately, the consequences of these interactions have not been convincingly correlated to patient report of fatigue. As well, findings of putative mechanism-directed therapies have not been successful to date. One possible mechanism currently being studied, however, involves the inflammatory response to cancer and its therapy, especially cytokine activation. Clinical studies have revealed increases in interleukin-1 (IL-1) receptor antagonist and C-reactive protein in fatigued patients with cancer who are undergoing radiation therapy. CFR has also been correlated with tumor necrosis factor (sTNF-RII), immunoregulatory CD4 lymphocytes, and white blood cell subsets. Proinflammatory proteins have been noted in breast cancer survivors with no evidence of disease who have experienced fatigue 3 to 5 years after cancer therapy. The reasons why inflammatory responses are still maintained after treatment and resolution of the cancer have not been fully explained. In a study of a group of breast cancer survivors with or without fatigue, at least one cytokine polymorphism at IL-1 β -511 ($P = .007$) was associated with reported fatigue.

Regardless of the quality or quantity of the pathophysiology involved, patients consistently note multiple sensations that often occur not in isolation but in clusters of complaints. Given the interwoven nature of the facets of CRF, causative perturbations may exert a systemic effect on multiple organs and tissues to obtain the phenotype of fatigue. Clinically, this cluster appears similar to inflammatory-induced symptoms referred to as “sickness behavior” and may encompass depression, sleep disturbances, and cognitive dysfunction. A sickness behavior animal model induced by lipopolysaccharide injection has been associated with behavioral changes (e.g., listlessness, decreased oral intake, and sleep disturbances) and increased levels of circulating proinflammatory cytokines. Although this preclinical model has been widely used for CRF studies, a lack of valid preclinical models of fatigue remains. The discovery of new animal models of fatigue continues to be a challenging yet key aspect of CRF research. Because CFR is subjective, the patients' description of fatigue (at the bedside) should guide the generation of hypotheses (at the bench) to begin to uncover the etiology or etiologies of CRF, even though some questions surrounding CRF mechanisms do not easily lend themselves to preclinical assessment help to accelerate an improved mechanistic understanding of fatigue. This understanding will allow for more precise interventions to be brought into clinical testing. Several grants and research projects are currently exploring inflammation as the primary driver of CRF.




FIG :4 Pathophysiology of Cancer

PATHOPHYSIOLOGY OF CANCER

The pathophysiology of cancer involves several stages. The first stage is initiation, where a mutation occurs in a cell's DNA, leading to the activation of oncogenes (genes that promote cell growth) or the inactivation of tumor suppressor genes (genes that inhibit cell growth). The second stage is promotion, where the mutated cells are stimulated to divide and grow rapidly, forming a small cluster of abnormal cells. The third stage is progression, where the abnormal cells continue to divide and grow, forming a tumor that can invade surrounding tissues and spread to other parts of the body through the bloodstream or lymphatic system.

One of the fundamental features of cancer is tumor clonality, the development of tumors from single cells that begin to proliferate abnormally. The single-cell origin of many tumors has been demonstrated by analysis of X chromosome inactivation. As discussed in Chapter 8, one member of the X chromosome pair is inactivated by being converted to heterochromatin in female cells. X inactivation occurs randomly during embryonic development, so one X chromosome is inactivated in some cells, while the other X chromosome is inactivated in other cells. Thus, if a female is heterozygous for an X chromosome gene, different alleles will be expressed in different cells. Normal tissues are composed of mixtures of cells with different inactive X chromosomes, so expression of both alleles is detected in normal tissues of heterozygous females. In contrast, tumor tissues generally express only one allele of a heterozygous X chromosome gene.

At the cellular level, the development of cancer is viewed as a multistep process involving mutation and selection for cells with progressively increasing capacity for proliferation, survival, invasion, and metastasis. The first step in the process, tumor initiation, is thought to be the result of a genetic alteration leading to abnormal proliferation of a single cell. Cell proliferation then leads to the outgrowth of a population of clonally derived tumor cells. Tumor progression continues as additional mutations occur within cells of the tumor population. Some of these mutations confer a selective advantage to the cell, such as more rapid growth, and

the descendants of a cell bearing such a mutation will consequently become dominant within the tumor population. The process is called clonal selection, since a new clone of tumor cells has evolved based on its increased growth rate or other properties (such as survival, invasion, or metastasis) that confer a selective advantage. Clonal selection continues throughout tumor development, so tumors continuously become more rapid-growing and increasingly malignant.

Studies of colon carcinomas have provided a clear example of tumor progression during the development of a common human malignancy. The earliest stage in tumor development is increased proliferation of colon epithelial cells. One of the cells within this proliferative cell population is then thought to give rise to a small benign neoplasm (an adenoma or polyp). Further rounds of clonal selection lead to the growth of adenomas of increasing size and proliferative potential. Malignant carcinomas then arise from the benign adenomas, indicated by invasion of the tumor cells through the basal lamina into underlying connective tissue. The cancer cells then continue to proliferate and spread through the connective tissues of the colon wall. Eventually the cancer cells penetrate the wall of the colon and invade other abdominal organs, such as the bladder or small intestine. In addition, the cancer cells invade blood and lymphatic vessels, allowing them to metastasize throughout the body.




FIG :5 Difference between normal cell and cancer cell

SIGNS AND SYMPTOMS OF CANCER

1. A persistent sore especially if it is in the mouth, on the lip or on tongue. Long standing sores due to smoking, sharp edge of teeth, or dental plates that do not fit, can develop into cancer if not taken care of. Common 'Cold Sore' disappears after a few days
2. Cancer does not. If any such sore persists more than a few days, see your doctor. Lumps or masses which appear where none were present previously should arouse suspicion of cancer.
3. This is especially true in the case of women and is the most readily cured if treated early. Persistent indigestion or change in bowel habits, while usually due to other causes, are often caused in the intestinal tract and should be reported.

RISK FACTORS

It is usually not possible to know exactly why one person develops cancer and another doesn't. But research has shown that certain risk factors may increase a person's chances of developing cancer. (There are also factors that are linked to a lower risk of cancer. These are sometimes called protective risk factors, or just protective factors.)

Cancer risk factors include exposure to chemicals or other substances, as well as certain behaviors. They also include things people cannot control, like age and family history.

SOME RISK FACTORS

Alcohol Cancer-Causing Substances

Chronic Inflammation

Diet Hormones

Immunosuppression Infectious Agents

Obesity

Radiation

DIAGNOSIS OF CANCER



The diagnosis of cancer involves several tests, including imaging tests such as X-rays, CT scans, and MRI scans, as well as laboratory tests such as biopsies and blood tests. Biopsies involve taking a small sample of tissue from the tumor and examining it under a microscope to determine if it is cancerous or benign (non-cancerous). Blood tests can detect certain substances in the blood that are associated with cancer, such as tumor markers.

1. Blood chemistry test

What it does: A blood chemistry test measures the amounts of certain substances that are released into the blood by the organs and tissues of the body. These substances include metabolites, electrolytes, fats, sugars, and proteins, including enzymes.

What it tells us: Blood chemistry tests give important information about how well your kidneys, liver, and other organs are working. High or low levels of some substances in the blood can be a sign of disease or treatment side effects.

2.Complete blood count

What it does: The CBC measures the number of red blood cells, white blood cells, and platelets in your blood. It also measures the amount of hemoglobin (the protein that carries oxygen) in your blood, the amount of your blood that is made up of red blood cells (hematocrit)

How it is used: The CBC is often part of a routine health check-up. It can help diagnose some cancers, especially leukemias. It is also used to monitor your health during and after treatment.in your red blood cell.

3.Cytogenetic

How it is used: The CBC is often part of a routine health check-up. It can help diagnose some cancers, especially leukemias. It is also used to monitor your health during and after treatment.

What it does: Cytogenetic analysis looks for changes in chromosomes in samples of tissue, blood, bone marrow, or amniotic fluid. Chromosome changes may include broken, missing, rearranged, or extra chromosomes. Changes in certain chromosomes may be a sign of a genetic condition or some types of cancer.

4.Immunophenotyping

What it does: Immunophenotyping uses antibodies to identify cells based on the types of antigens or markers on the surface of the cells. It is most often done on blood or bone marrow samples. But it may also be done on other body fluids or tissue samples.

How it is used: Immunophenotyping helps diagnose, stage, and monitor blood cancers and other blood disorders, such as leukemias, lymphomas, myelodysplastic syndromes,

TREATMENT OF CANCER

Alternative cancer treatment; Integrative medicine treatments are meant to be used along with your cancer treatments. Integrative treatments generally aren't powerful enough to replace standard treatments. They may help you cope with symptoms caused by cancer and cancer treatments, such as anxiety, fatigue, nausea and vomiting, pain, problems sleeping, and stress. Talk about your options with your healthcare team to help find

FEW TREATMENTS.ARE LIKE

1.Accupuncture

During acupuncture treatment, an acupuncture specialist inserts tiny needles into your skin at precise points. Studies show acupuncture may be helpful in improving sleep and reducing nausea caused by chemotherapy. Acupuncture also may help relieve certain types of pain in people with cancer. It may help with hot flashes in those with breast cancer who are taking medicines that may affect hormone levels.

2.Aromatherapy

Aromatherapy uses fragrant oils to provide a calming sensation. Oils infused with scents such as lavender can be applied to your skin during a massage. You also can add the oils to bath water. Fragrant oils can be heated to release their scents into the air. Aromatherapy may be helpful in relieving nausea. While there are trained professionals who offer aromatherapy, you also may use aromatherapy on your own. Aromatherapy is safe, although oils applied to the skin can cause allergic reactions. People with cancer that is sensitive to the effects of estrogen, such as some breast cancers, may want to avoid applying large amounts of lavender oil or tea tree oil to the skin. These oils may have effects like hormones if you apply too much.

3.Hypnosis

Hypnosis is a state of deep concentration. During a hypnotherapy session, a therapist may hypnotize you by talking in a gentle voice and helping you relax. The therapist helps you focus on goals, such as less pain. Hypnosis may be helpful for people with cancer who are having pain and stress. It may be most helpful for people having surgery or procedures to help treat their cancer. When performed by a certified therapist, hypnosis is safe. But tell your therapist if you have a history of mental illness.

4.Meditation

Meditation is a state of deep concentration when you focus your mind on one image, sound or idea, such as a positive thought. When meditating, you also might do deep-breathing or relaxation exercises. Meditation may help people with cancer by relieving anxiety and stress and improving mood. Meditation is generally safe. You can meditate on your own for a few minutes once or twice a day. You also can take a class with an instructor.

5.Music therapy

During music therapy sessions, you might listen to music, play instruments, sing songs or write lyrics. A trained music therapist may lead you through activities designed to meet your specific needs, or you may take part in music therapy in a group setting. Music therapy may help relieve pain and help you deal with anxiety and stress. Music therapy is safe, and participation doesn't require any musical talent or training. Many medical centers have certified music therapists on staff.

6.Yoga

Yoga combines stretching exercises with deep breathing. During a yoga session, you position your body in various poses that require bending, twisting and stretching. There are many types of yoga, each with its own movements. Yoga may provide some stress relief for people with cancer. Yoga has also been shown to improve sleep and reduce fatigue.

7.Massage.

During a massage, a trained professional kneads your skin, muscles and tendons to try to relieve muscle tension and stress and promote relaxation. Several massage methods exist. Massage can be light and gentle, or it can be deep with more pressure. Studies have found that massage can be helpful in relieving pain in people with cancer. It also may help relieve fatigue and stress. Massage can be safe if you work with a professional massage therapist. Many cancer centers have massage therapists on staff. Your healthcare professional may refer you to a massage therapist who often works with people who have cancer. Avoid getting a massage if your blood counts are very low. Ask the massage therapist to avoid massaging near surgical scars, radiation treatment areas or tumors. If you have cancer in your bones or have other bone diseases, such as osteoporosis, ask the massage therapist to use light pressure rather than deep massage.

8.Tai chi.

Tai chi is a form of exercise that includes gentle movements and deep breathing. Tai chi can be led by an instructor, or you can learn tai chi on your own following books or videos. Practicing tai chi may help relieve stress and lower fatigue. Tai chi is generally safe. The slow movements of tai chi don't require great physical strength, and you can usually adapt the exercises to fit how well you move. Still, talk to a member of your healthcare team before beginning tai chi. Avoid doing any tai chi moves that cause pain

Conclusion on Management of Stage 1

Stage 1 cancer represents a critical window where early detection and timely intervention can lead to significantly better outcomes. The document highlights that while conventional Cancer Without Medication treatments like surgery, chemotherapy, and radiation remain the backbone of cancer management, non-pharmacological and integrative approaches—such as meditation, yoga, hypnosis, acupuncture, aromatherapy, music therapy, massage, and tai chi—play an important supportive role.

These therapies do not replace medical treatment but can:

1. Reduce stress, anxiety, and fatigue
2. Improve mood, sleep, and overall quality of life
3. Help patients cope with pain and treatment side effects
4. Encourage holistic well-being through mind–body balance

Ultimately, the management of Stage 1 cancer without medication emphasizes early detection, lifestyle modification, and supportive therapies that enhance resilience and recovery. By combining these approaches with medical guidance, patients can experience fewer adverse effects and maintain a better quality of life during their cancer journey.

REFERENCE

- [1] Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021; 71: 209–249.
- [2] Ganesh K, Massagué J. Targeting metastatic cancer. *Nat Med* 2021; 27: 34–44.
- [3] Johnson, S. B., Park, H. S., Gross, C. P., & Yu, J. B. (2018). Use of alternative medicine for cancer. *Journal of the National Cancer Institute*, 110:121-124.
- [4] Chow, R., Zhao, H., DeAngelis, C., et al. (2020) Integrative Medicine in Oncology. *Journal of Supportive Care in Cancer*, 28: 4021-4030.
- [5] Cheng K, Wong W, Koh C (2016) Unmet Needs Mediate the Relationship Between Symptoms and Quality of Life in Breast Cancer Survivors. *Supportive Care Cancer* 24(5).
- [6] Palesh O, Scheiber C, Kesler S, Mustian K, Koopman C, Schapira L (2018) Management of Side Effects During and Post-Treatment in Breast Cancer Survivors. *Breast J* 24(2).
- [7] Horton S, Camacho Rodriguez R, Anderson BO, Aung S, Awuah B, Delgado Pebe L, et al (2020) Health System Strengthening: Integration of Breast Cancer Care for Improved Outcomes. *Cancer* 126:2353–64.
- [8] Foster C, Calman L, Richardson A, Pimperton H, Nash R (2018) Improving the Lives of People Living with and Beyond Cancer: Generating the Evidence Needed to Inform Policy and Practice. *J Cancer Policy* 15:92–5.
- [9] Halpern MT, Viswanathan M, Evans TS, Birken SA, Basch E, Mayer DK (2015) Models of Cancer Survivorship Care: Overview and Summary of Current Evidence. *J Oncol Practice* 11(1):e19–27.
- [10] Shneerson C, Taskila T, Holder R, Greenfield S, Tolosa I, Damery S, et al (2015) Patterns of Self-Management Practices Undertaken by Cancer Survivors: Variations in Demographic Factors. *Eur J Cancer Care* 24(5).

- [11] Cuthbert CA, Farragher JF, Hemmelgarn BR, Ding Q, McKinnon GP, Cheung WY (2019) Self-Management Interventions for Cancer Survivors: A Systematic Review and Evaluation of Intervention Content and Theories. *Psycho-Oncol* 28(11).
- [12] Boland L, Bennett K, Connolly D (2018) Self-Management Interventions for Cancer Survivors: A Systematic Review. *Supportive Care Cancer* 26(5).
- [13] Richardson J, Loyola-Sanchez A, Sinclair S, Harris J, Letts L, MacIntyre NJ, et al (2014) Self-Management Interventions for Chronic Disease: A Systematic Scoping Review. *Clin Rehabilitation* 28(11).
- [14] Hammer MJ, Ercolano EA, Wright F, Dickson VV, Chyun D, Melkus GDE (2015) Self-Management for Adult Patients with Cancer: An Integrative Review. *Cancer Nurs* 38(2).

Copyright & License:

© Authors retain the copyright of this article. This work is published under the Creative Commons Attribution 4.0 International License (CC BY 4.0), permitting unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.